

## Catheters & Ports in Cancer Treatment

### What are catheters and ports?

Catheters are long, narrow, hollow tubes made of soft plastic that are inserted into a vein. Your health care team uses them to give medications, blood transfusions, or fluids or to take samples of blood for testing. Treatments are given through a small device connected to the catheter. Your doctor may put a catheter completely under the skin. If so, it is connected to a small plastic or metal disc called a port. The entire device is called a port-o-cath. A port may remain in place for years, if necessary.



### What are the types of catheters?

The peripheral venous catheter (PVC) is the most common type. It is a short catheter, typically inserted into a vein in the arm or hand. It stays in place no longer than 3 days.

A peripherally inserted central catheter (PICC) is inserted into a large vein near your elbow. A central line is inserted

into a large vein under the collarbone or in the neck. The tube goes under the skin from where it was inserted to where it leaves the body, usually in the upper chest. The exit is where your doctor gives you medication. A PICC line or CVC may remain in place for weeks or months.

### What are the benefits of a catheter or port?

Both catheters and ports reduce the number of times you need to be stuck with a needle and allow more than 1 type of treatment or medication to be given at a time. Catheters also decrease the risk of tissue and muscle damage that can occur if some types of chemotherapy leak outside a vein, prevent excessive bruising or bleeding in patients with bleeding problems, and allow some chemotherapy treatments to be given at home.

### How should I take care of a catheter or port?

Taking good care of a catheter or port reduces the risk of developing side effects such as infections, blockages or clots, kinks under the skin, and movement of the catheter. For catheters with tips that remain outside the body, you must take special care of the tube and the skin surrounding the area where the tube exits the body. In addition, these catheters must be flushed with sterile fluid each day to prevent blockage. An IV care service can help with this until you are comfortable doing it yourself. Ports require less care because they are located completely under the skin. A port may still need to be flushed so it does not get blocked. Ask your health care team for specific instructions about how to care for your catheter or port.

### What are the signs of problems with a catheter or port?

Contact your doctor immediately if the area around the catheter or port becomes red, swollen, painful, bruised, or warm; there is excess bleeding from the insertion area; you develop a fever; you have shortness of breath or dizziness; fluid leaks; the catheter tube outside of your body gets longer; or the catheter seems blocked.

## Questions to ask the doctor

Regular communication is important for making informed decisions about your health care. Consider asking the following questions of your health care team:

- Why are you recommending a catheter or port?
- What are the risks and benefits of a catheter or port?
- Will my health insurance cover the costs associated with inserting a catheter or port?
- Will I receive anesthesia when the catheter or port is inserted? If so, which type?
- How long will the procedure take? How long will I need to stay in the hospital?
- How long will the catheter or port be left in place?
- How should I care for my catheter or port?
- May I wear regular clothes, bathe or swim, or exercise with my catheter or port?
- What are signs of a problem with my catheter or port?
- Whom should I contact if I have problems with my catheter or port?
- Is there anything else I should be asking?

Find additional cancer treatment information at [www.cancer.net/treatment](http://www.cancer.net/treatment).

---

The ideas and opinions expressed here do not necessarily reflect the opinions of the American Society of Clinical Oncology (ASCO) or The Conquer Cancer Foundation. The information in this fact sheet is not intended as medical or legal advice, or as a substitute for consultation with a physician or other licensed health care provider. Patients with health care-related questions should call or see their physician or other health care provider promptly and should not disregard professional medical advice, or delay seeking it, because of information encountered here. The mention of any product, service, or treatment in this fact sheet should not be construed as an ASCO endorsement. ASCO is not responsible for any injury or damage to persons or property arising out of or related to any use of ASCO's patient education materials, or to any errors or omissions.

To order more printed copies, please call 888-273-3508 or visit [www.cancer.net/estore](http://www.cancer.net/estore).



### AMERICAN SOCIETY OF CLINICAL ONCOLOGY

2318 Mill Road, Suite 800, Alexandria, VA 22314 | Toll Free: 888-651-3038 | Phone: 571-483-1300

[www.asco.org](http://www.asco.org) | [www.cancer.net](http://www.cancer.net) | [www.conquercancerfoundation.org](http://www.conquercancerfoundation.org)

© 2016 American Society of Clinical Oncology. For permissions information, contact [permissions@asco.org](mailto:permissions@asco.org).

## TERMS TO KNOW

### Anesthesia:

Medication to block the awareness of pain

### Bolus:

A dose of a substance given intravenously; also called an IV push

### Chemotherapy:

The use of drugs to destroy cancer cells

### Continuous infusion:

Intravenous treatment delivered over a specific time without interruption

### Displacement:

A shift in the position of a catheter

### Drip chamber:

A device that regulates the flow of substances given intravenously and prevents air from entering the bloodstream

### Infiltration:

A complication when intravenous fluid or medication enters the tissue surrounding the catheter, rather than entering the vein; also called extravasation

### Infusion:

The process of delivering a substance into a vein

### Infusion pump:

A medical device that delivers treatment intravenously in controlled amounts

### Intermittent infusion:

Intravenous treatment delivered only at certain times and used when a person does not need additional fluid

### Phlebitis:

Inflammation of a vein

MADE AVAILABLE THROUGH

